




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>C07D 213/75, 215/38, 215/46, A61K</b> <b>31/47, 31/44</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 94/18170</b> <b>(43) International Publication Date:</b> 18 August 1994 (18.08.94)
<b>(21) International Application Number:</b> PCT/EP94/00189 <b>(22) International Filing Date:</b> 25 January 1994 (25.01.94)  <b>(30) Priority Data:</b> 9302275.4      5 February 1993 (05.02.93)      GB  <b>(71) Applicant (for all designated States except US):</b> SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex 9EP TW8 (GB).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> FORBES, Ian, Thomson [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). HAM, Peter [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). MARTIN, Roger, Thomas [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). THOMPSON, Mervyn [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB).		<b>(74) Agent:</b> GIDDINGS, Peter, J.; Corporate Intellectual Property, SmithKline Beecham, Mundells, Welwyn Garden City, Hertfordshire AL7 1EY (GB).  <b>(81) Designated States:</b> JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>

**(54) Title:** USE OF PHENYL HETEROARYL UREAS AS 5HT<sub>2c</sub> RECEPTOR ANTAGONISTS AND UREA COMPOUNDS

**(57) Abstract**

The use of a compound of formula (I) or a salt thereof, wherein P represents a quinoline or isoquinoline residue or a 5- or 6-membered aromatic heterocyclic ring containing up to three heteroatoms selected from nitrogen, oxygen or sulphur; R<sup>1</sup> is hydrogen, C<sub>1-6</sub> alkyl, halogen, NR<sup>5</sup>R<sup>6</sup> or OR<sup>7</sup> where R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are independently hydrogen or C<sub>1-6</sub> alkyl; R<sup>2</sup> and R<sup>3</sup> are independently hydrogen or C<sub>1-6</sub> alkyl; R<sup>4</sup> is hydrogen, C<sub>1-6</sub> alkyl, CF<sub>3</sub>, nitro, cyano, acyl, halogen, NR<sup>5</sup>R<sup>6</sup>, OR<sup>7</sup> or CO<sub>2</sub>R<sup>7</sup> where R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are independently hydrogen or C<sub>1-6</sub> alkyl as defined for R<sup>1</sup>; and n is 1, 2 or 3, in the manufacture of a medicament for the treatment or prophylaxis of CNS disorders.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				


# USE OF PHENYL HETEROARYL UREAS AS 5HT<sub>2C</sub> RECEPTOR ANTAGONISTS AND UREA COMPOUNDS

This invention relates to a method of treatment of certain CNS disorders.

- 5 WO 92/05170 describes certain urea derivatives which are described as possessing 5HT<sub>1C</sub> receptor antagonist activity. The 5HT<sub>1C</sub> receptor has recently been reclassified as the 5HT<sub>2C</sub> receptor [P. Hartig et al., Trends in Pharmacological Sciences (TIPS) 1993].

- 10 Certain phenyl heteroaryl ureas known in the art have now been found to have 5HT<sub>2C</sub> receptor antagonist activity. 5HT<sub>2C</sub> receptor antagonists are believed to be of potential use in the treatment of CNS disorders such as anxiety, depression, obsessive compulsive disorders, migraine, anorexia, Alzheimers disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse such as cocaine, ethanol, nicotine and benzodiazepines, schizophrenia, and also disorders associated with spinal trauma and/or head injury such as
- 15 hydrocephalus.

Accordingly, the present invention provides the use of a compound of formula (I) or a salt thereof:



20

wherein:

- P represents a quinoline or isoquinoline residue or a 5- or 6-membered aromatic heterocyclic ring containing up to three heteroatoms selected from nitrogen, oxygen or
- 25 sulphur;
- R<sup>1</sup> is hydrogen, C<sub>1-6</sub> alkyl, halogen, NR<sup>5</sup>R<sup>6</sup> or OR<sup>7</sup> where R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are independently hydrogen or C<sub>1-6</sub> alkyl;
- R<sup>2</sup> and R<sup>3</sup> are independently hydrogen or C<sub>1-6</sub> alkyl;
- R<sup>4</sup> is hydrogen, C<sub>1-6</sub> alkyl, CF<sub>3</sub>, nitro, cyano, acyl, halogen, NR<sup>5</sup>R<sup>6</sup>, OR<sup>7</sup> or CO<sub>2</sub>R<sup>7</sup>
- 30 where R<sup>5</sup>, R<sup>6</sup> and R<sup>7</sup> are independently hydrogen or C<sub>1-6</sub> alkyl as defined for R<sup>1</sup>; and n is 1, 2 or 3,
- in the manufacture of a medicament for the treatment or prophylaxis of CNS disorders.

C<sub>1-6</sub>alkyl groups, whether alone or as part of another group, can be straight chain or branched.

- 5 Preferably R<sup>1</sup> is hydrogen or methyl.

Preferably R<sup>2</sup> and R<sup>3</sup> are hydrogen.

- 10 Suitable moieties when the ring P is a 5- or 6-membered aromatic heterocyclic ring include pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, isothiazolyl, isoxazolyl, thiadiazolyl and triazolyl. Preferably P is pyridyl attached to the urea nitrogen at position 3 or 4; or P is quinoline attached to the urea nitrogen at position 3, 4 or 6, preferably at position 4.

- 15 Preferably n is 1 or 2. When n is greater than 1, the R<sup>4</sup> groups can be the same or different. Preferably the phenyl ring is mono-substituted and R<sup>4</sup> is CF<sub>3</sub> or -NMe<sub>2</sub> (preferably in the meta position); -OMe, (preferably in the meta or para position); CO<sub>2</sub>Et (preferably in the meta position) or the phenyl ring is preferably di substituted with meta chloro and para methyl.

- 20 Preferred compounds of formula (I) include:  
 N-(Phenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(6-Quinoliny)l)-N'-(3-trifluoromethylphenyl) urea,  
 N-(3-Dimethylaminophenyl)-N'-(6-quinoliny)l urea,  
 N-(Phenyl)-N'-(6-quinoliny)l urea,  
 25 N-(4-Methoxyphenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(3-Dimethylaminophenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(3-Methoxyphenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(3-Ethoxycarbonylphenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(2-Methyl-4-quinoliny)l)-N'-(3-trifluoromethylphenyl) urea ,  
 30 N-(Phenyl)-N'-(3-quinoliny)l urea,  
 N-(3-Chloro-4-methylphenyl)-N'-(3-pyridyl) urea,  
 N-(3-Chloro-4-methylphenyl)-N'-(4-pyridyl) urea,  
 N-(3-Pyridyl)-N'-(3-(trifluoromethyl)phenyl)urea,  
 N-(3-Methylphenyl)-N'-(3-pyridyl)urea,  
 35 N-(4-Chlorophenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chlorophenyl)-N'-(3-pyridyl)urea,  
 N-(3-Hydroxyphenyl)-N'-(2-methyl-4-quinoliny)lurea,  
 N-(3-Bromophenyl)-N'-(3-pyridyl)urea.

- N-(3,4-Dichlorophenyl)-N'-(3-pyridyl)urea,  
N-(3-Fluoro-4-methylphenyl)-N'-(3-pyridyl)urea,  
N-(4-Ethoxycarbonylphenyl)-N'-(3-pyridyl)urea,  
N-(3-Chloro-4-methoxycarbonylphenyl)-N'-(3-pyridyl)urea,  
5 N-(3-Bromo-4-methylphenyl)-N'-(3-pyridyl)urea,  
N-(3-Chloro-4-cyanophenyl)-N'-(3-pyridyl)urea,  
N-(4-Nitro-3-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
N-(4-Chloro-3-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
N-(3-Chloro-4-carboxyphenyl)-N'-(3-pyridyl)urea,  
10 N-(2-Methoxy-4-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
N-(3-Chloro-4-ethylphenyl)-N'-(3-pyridyl)urea,  
N-(3-Chloro-4-propylphenyl)-N'-(3-pyridyl)urea,  
N-(3-Chloro-4-tert-butylphenyl)-N'-(3-pyridyl)urea,  
N-(3-Hydroxy-4-(methoxycarbonyl)phenyl)-N'-(3-pyridyl)urea  
15 or a pharmaceutically acceptable salt thereof.

- The compounds of the formula (I) can form acid addition salts with acids, such as conventional pharmaceutically acceptable acids, for example maleic, hydrochloric, hydrobromic, phosphoric, acetic, fumaric, salicylic, citric, lactic, mandelic, tartaric and  
20 methanesulphonic. Compounds of formula (I) may also form N-oxides or solvates such as hydrates, and the invention also extends to these forms.

- Certain compounds of formula (I) may exist tautomerically in more than one form. The invention extends to these and any other tautomeric forms and mixtures thereof.  
25 Certain compounds of formula (I) are capable of existing in stereoisomeric forms including enantiomers and the invention extends to each of these stereoisomeric forms and to mixtures thereof including racemates. The different stereoisomeric forms may be separated one from the other by the usual methods, or any given isomer may be obtained by stereospecific or asymmetric synthesis.

- 30 Certain compounds of formula (I) are novel and form a further aspect of the invention. Particularly preferred novel compounds include those listed above and exemplified herein.

- The invention further provides a method of treatment or prophylaxis of CNS disorders, in  
35 particular anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse, schizophrenia and/or disorders associated with spinal trauma and/or head injuries (in particular anxiety and depression) in mammals including humans, which comprises

administering to the sufferer a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

5 The invention also provides novel compounds of formula (I) or a pharmaceutically acceptable salt thereof, for use as a therapeutic substance, in particular in the treatment or prophylaxis of anxiety, depression, migraine, anorexia, obsessive compulsive disorders, Alzheimer's disease, sleep disorders, bulimia, panic attacks, withdrawal from drug abuse, schizophrenia and/or disorders associated with spinal trauma and/or head injuries.

10 The present invention also provides a pharmaceutical composition, which comprises novel compounds of formula (I) or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

15 A pharmaceutical composition of the invention, which may be prepared by admixture, suitably at ambient temperature and atmospheric pressure, is usually adapted for oral, parenteral or rectal administration and, as such, may be in the form of tablets, capsules, oral liquid preparations, powders, granules, lozenges, reconstitutable powders, injectable or infusable solutions or suspensions or suppositories. Orally administrable compositions are generally preferred.

20 Tablets and capsules for oral administration may be in unit dose form, and may contain conventional excipients, such as binding agents, fillers, tableting lubricants, disintegrants and acceptable wetting agents. The tablets may be coated according to methods well known in normal pharmaceutical practice.

25 Oral liquid preparations may be in the form of, for example, aqueous or oily suspension, solutions, emulsions, syrups or elixirs, or may be in the form of a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, emulsifying agents, 30 non-aqueous vehicles (which may include edible oils), preservatives, and, if desired, conventional flavourings or colourants.

For parenteral administration, fluid unit dosage forms are prepared utilising a compound of the invention or pharmaceutically acceptable salt thereof and a sterile vehicle. The 35 compound, depending on the vehicle and concentration used, can be either suspended or dissolved in the vehicle. In preparing solutions, the compound can be dissolved for injection and filter sterilised before filling into a suitable vial or ampoule and sealing. Advantageously, adjuvants such as a local anaesthetic, preservatives and buffering agents

are dissolved in the vehicle. To enhance the stability, the composition can be frozen after filling into the vial and the water removed under vacuum. Parenteral suspensions are prepared in substantially the same manner, except that the compound is suspended in the vehicle instead of being dissolved, and sterilization cannot be accomplished by filtration.


- 5 The compound can be sterilised by exposure to ethylene oxide before suspension in a sterile vehicle. Advantageously, a surfactant or wetting agent is included in the composition to facilitate uniform distribution of the compound.

- 10 The composition may contain from 0.1% to 99% by weight, preferably from 10 to 60% by weight, of the active material, depending on the method of administration.


- The dose of the compound used in the treatment of the aforementioned disorders will vary in the usual way with the seriousness of the disorders, the weight of the sufferer, and other similar factors. However, as a general guide suitable unit doses may be 0.05 to 1000 mg,  
15 more suitably 0.05 to 20.0 mg, for example 0.2 to 5 mg; and such unit doses may be administered more than once a day, for example two or three a day, so that the total daily dosage is in the range of about 0.01 to 100 mg/kg; and such therapy may extend for a number of weeks or months.

- 20 When administered in accordance with the invention, no unacceptable toxicological effects are expected with the compounds of the invention.

- The present invention also provides a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof, which process comprises  
25 the coupling of a compound of formula (II);



- 30 with a compound of formula (III);



- 35 wherein P is as defined in relation to formula (I), A and B contain the appropriate functional group(s) necessary to form the moiety,  $-NR^{2'}CONR^{3'}$  when coupled, the

- variables  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  are  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  respectively, as defined in formula (I), or groups convertible thereto, and thereafter optionally and as necessary and in any appropriate order, converting any  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$ , when other than  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  respectively to  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$ , interconverting  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  and
- 5 forming a pharmaceutically acceptable salt thereof.

Suitable examples of groups A and B include:

- (i) A is  $-N=C=O$  and B is  $-NHR^{3'}$ ,
- 10 (ii) A is  $-NR^{2'}COL$  and B is  $-NHR^{3'}$ ,
- (iii) A is  $-NHR^{2'}$  and B is  $NR^{3'}COL$ ,
- (iv) A is  $NHR^{2'}$  and B is  $-N=C=O$  or
- (v) A is halogen and B is  $-NR^{3'}CONHR^{2'}$

- 15 wherein  $R^{2'}$  and  $R^{3'}$  are as defined above and L is a leaving group. Examples of suitable leaving groups L include halogen such as chloro, bromo, imidazole or phenoxy or phenylthio optionally substituted for example with halogen.

- When A is  $-N=C=O$  and B is  $NHR^{3'}$  or when A is  $NHR^{2'}$  and B is  $-N=C=O$  the reaction is
- 20 suitably carried out in an inert solvent for example dichloromethane or toluene at ambient temperature. When A is  $-NR^{2'}COL$  and B is  $NHR^{3'}$  or when A is  $-NHR^{2'}$  and B is  $-NR^{3'}COL$ , the reaction is suitably carried out in an inert solvent such as dichloromethane at ambient temperature optionally in the presence of a base, such as triethylamine or in dimethylformamide at ambient or elevated temperature. When A is halogen and B is
- 25  $NR^{3'}CONHR^{2'}$ , the reaction is suitably carried out in an inert solvent such as toluene at elevated temperature, optionally in the presence of a base.

- Suitable examples of groups  $R^1$  and  $R^4$ , which are convertible to  $R^1$  and  $R^4$  alkyl groups respectively, include acyl groups which are introduced conventionally and may be
- 30 converted to the corresponding alkyl group by conventional reduction, such as using sodium borohydride in an inert solvent followed by hydrogenolysis in an inert solvent. Hydrogen substituents may be obtained from alkoxycarbonyl groups which may be converted to hydrogen by hydrolysis and decarboxylation.

- 35 Interconversions of  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  are carried out by conventional procedures. For example, in the case wherein  $R^2$  is  $C_{1-6}$  alkyl and  $R^3$  is hydrogen it is possible to introduce a  $C_{1-6}$  alkyl group at the  $R^3$  position by conventional alkylation using 1 molar equivalent of a  $C_{1-6}$  alkyl halide and 1 molar equivalent of a suitable base in an inert solvent. Suitable examples of a group  $R^{2'}$  and  $R^{3'}$  which is convertible to  $R^2$  and  $R^3$ ,



include alkoxycarbonyl and benzyl or *para*-methoxybenzyl which are converted to R<sup>2</sup> and R<sup>3</sup> is hydrogen using conventional conditions.

5 R<sup>1</sup> halo and R<sup>4</sup> halo may be introduced by selective halogenation of the ring P or the benzene ring respectively using conventional conditions.

It should be appreciated that it may be necessary to protect any R<sup>1</sup> to R<sup>7</sup> hydrogen variables which are not required to be interconverted. Suitable protecting groups are described in 'Protective groups in organic synthesis' Greene T.W., New York, Wiley  
10 (1981). It should be appreciated that it is preferred that groups R<sup>1</sup> to R<sup>7</sup> are introduced before coupling compounds of formula (II) and (III).

Compounds of formula (II) in which A is NHR<sup>2'</sup> are known compounds or can be prepared analogously to known compounds. see, for example, WO 92/05170 (SmithKline  
15 Beecham plc). Compounds of formula (II) in which A is -N=C=O may be prepared by treating a compound of formula (II) in which :

- i) A is amino, with phosgene or a phosgene equivalent, in the presence of excess base in an inert solvent.
- 20 ii) A is acylazide (i.e. CON<sub>3</sub>), via the nitrene, by thermal rearrangement using conventional conditions (ref L.S. Trifonov et al, Helv. Chim. Acta 1987 70 262).
- iii) A is CONH<sub>2</sub>, via the nitrene intermediate using conventional conditions.

Examples of phosgene equivalents include triphosgene, carbonyldiimidazole, phenyl  
25 chloroformate and phenyl chlorothioformate. Compounds of formula (II) in which A is NR<sub>2'</sub>COL may be prepared by reacting a compound of formula (II) in which A is NHR<sub>2'</sub> with phosgene or a phosgene equivalent in an inert solvent, at low temperature, if necessary in the presence of one equivalent of a base such as triethylamine. Compounds of formula (II) in which A is halogen and R<sub>4'</sub> is hydrogen are commercially available.

30 Compounds of formula (III) are commercially available or may be prepared according to analogous methods to those outlined above for compounds of formula (II).

Pharmaceutically acceptable salts may be prepared conventionally by reaction with the  
35 appropriate acid or acid derivative. N-oxides may be formed conventionally by reaction with hydrogen peroxide or percarboxylic acids.

The following Examples illustrate the preparation of compounds of the invention.

**Example 1****N-(Phenyl)-N'-(2-methyl-4-quinolinyl) urea**

To a solution of 2-methyl-4-amino-quinoline (1.58g; 10.0mM) in ethanol free chloroform  
5 (70 ml) at room temperature was added dropwise phenyl isocyanate (1.4 ml, 12.0 mM) in  
dry toluene (25 ml) over a period of 5 minutes. The whole was then stirred at room  
temperature for 18h followed by heating under reflux for 2h. After cooling to room  
temperature ethanol (20 ml) was added and after an additional 1h the solvent was removed  
10 under reduced pressure to give an oil. The oil was treated with ether (50 ml) to give an off  
white solid which was crystallised from ethyl acetate to give the title compound (E1)  
(1.37g, 60%) as a white solid. m.p. 201-3°C.

NMR (D<sub>6</sub>-DMSO) δ: 2.10 (3H,s), 7.05 (1H,t,J=7Hz), 7.36 (2H,t, J=7Hz), 7.47-7.64  
(3H,m), 7.70 (1H,t,J=7Hz), 7.89 (1H,d,J=7Hz), 8.08-8.20 (2H,m), 9.15 (1H, broad s),  
15 9.39 (1H, broad s).

Found: C, 73.74; H, 5.44; N, 15.29%

C<sub>17</sub>H<sub>15</sub>N<sub>3</sub>O requires: C, 73.63; H, 5.45; N, 15.15%

**Example 2****N-(6-Quinolinyl)-N'-(3-trifluoromethylphenyl) urea hydrochloride**

The title compound was prepared in 87% yield from 3-trifluoromethyl phenyl isocyanate  
and 6-aminoquinoline following a procedure similar to that in Example 1. Free base  
precipitated from the reaction mixture within a total of 3h at room temperature and was  
25 converted to the hydrochloride salt using hydrogen chloride in ether/ethanol.  
m.p. 208-13°C

NMR (D<sub>6</sub>-DMSO) δ: 7.33-7.50 (1H,m), 7.54-7.85 (2H,m), 7.95-8.28 (3H,m), 8.49  
(1H,d,J=8Hz), 8.60-8.75 (1H,m), 9.05-9.30 (2H,m), 10.19 (1H,s). 10.40 (1H,s).

30 Found: M<sup>+</sup> 331.0917 C<sub>17</sub>H<sub>12</sub>F<sub>3</sub>N<sub>3</sub>O requires 331.0964

**Example 3****N-(3-Dimethylaminophenyl)-N'-(6-quinolinyl) urea dihydrochloride**

35 To a solution of carbonyl diimidazole (0.40g, 2.47 mM) in dichloromethane (5 ml) was  
added 3-dimethylamino aniline (0.37g, 2.26 mM) in dichloromethane (5 ml). After  
stirring at room temperature for 0.5h, the solution was evaporated to dryness. The residue  
was taken up in dimethylformamide (5ml) and to this solution was added 6-amino

- quinoline (0.36g, 2.50 mM) in dimethylformamide (5 ml). The reaction mixture was heated to 90°C for 1h, then cooled and added dropwise to water with vigorous stirring. Later the solution was collected to give the crude product (0.65g) which was converted to the hydrochloride salt (E3) using hydrogen chloride in ether/ethanol (0.29g, 36%).
- 5 m.p. 185-90°C.
- NMR (D<sub>6</sub>-DMSO) δ: 3.09 (6H,s), 7.02-7.20 (1H,m), 7.21-7.49 (2H,m), 7.71 (1H, broad s), 7.90-8.15 (2H,m), 8.33 (1H,d,J=9Hz), 8.51-8.61 (1H,m), 9.02-9.18 (2H,m), 9.90 (1H, broad s), 10.39 (1H, broad s).
- 10 Found: 306.1465 C<sub>18</sub>H<sub>18</sub>N<sub>4</sub>O requires: 306.1516

#### Example 4

##### N-(phenyl)-N'-(6-quinolinyl) urea

- 15 The title compound (E4) was prepared in 56% yield from 6-aminoquinoline and phenylisocyanate following a procedure similar to that in Example 1. Free base precipitated from the reaction mixture and was recrystallised from ethanol.
- NMR (D<sub>6</sub>-DMSO) δ: 7.0 (1H,t,6Hz), 7.30 (2H,t,6Hz), 7.42-7.53 (3H,m), 7.71 (1H, dd, 6Hz,3Hz), 7.94 (1H,d,J=6Hz), 8.19 (1H,d,j=3Hz), 8.24-8.28 (1H,m), 8.71-8.77 (1H,m), 8.82 (1H,s), 9.08 (1H,s).
- 20

#### Example 5

##### N-(4-Methoxyphenyl)-N'-(2-methyl-4-quinolinyl) urea

- 25 4-Methoxyphenyl isocyanate (0.82 ml, 6.3 mmol), in dry dichloromethane (30 ml) was added slowly to 4-aminoquinoline (1 g, 6.3 mmol) in dry toluene (30 ml) under a nitrogen atmosphere. This was refluxed for 2.5h, then left at room temperature for 19h. The precipitate which formed was filtered off, washed with cold 1:1
- 30 toluene/dichloromethane and dried *in vacuo*. The crude product was chromatographed on silica gel using dichloromethane as the eluant to give the title compound (1.31 g, 68%) as a white solid. m.p. 180-185°C.
- NMR (D<sub>6</sub>-DMSO) δ: 2.59 (3H, s), 3.75 (3H, s), 6.93 (2H, m), 7.44 (2H, m), 7.6 (1H, t, J 6), 7.71 (1H, t, J 6), 7.89 (1H, d, J 8), 8.13 (2H, m), 9.11 (1H, s), 9.17 (1H, s)
- 35

Found: C, 70.32; H, 5.67; N, 13.44%

$C_{18}H_{17}N_3O_2$  requires C, 70.34; H, 5.58; N, 13.67%

Found:  $M^+$  307  $C_{18}H_{17}N_3O_2$  requires 307

## 5 Example 6

### N-(3-Dimethylaminophenyl)-N'-(2-methyl-4-quinolinyl) urea

1,1'-Carbonyldiimidazole (0.26g, 1.6 mmol), was added portionwise to a solution of 4-aminoquinaldine (0.23g, 1.47 mmol) in dry dichloromethane (15 ml), under a nitrogen atmosphere. After 1/2h, at room temperature, the solvent was evaporated off and the residue was taken up in DMF (5 ml). After addition of 3-(dimethylamino)aniline (0.2g, 1.47 mmol) in DMF (10ml), the reaction mixture was heated at 90°C for 1h. Addition of water after cooling to room temperature, gave the crude product as a precipitate, which was collected and dried *in vacuo*. Purification by column chromatography on silica gel, using dichloromethane as eluant gave the title compound (0.16g, 34%) as a light brown solid, m.p. 167-171°C.

NMR ( $D_6$ -DMSO)  $\delta$ : 2.6 (3H, s), 2.91 (6H, s), 6.42 (1H, m), 6.77 (1H, m), 6.98 (1H, s), 7.12 (1H, t, J 6), 7.59 (1H, t, J 3), 7.72 (1H, t, J 6), 7.89 (1H, d, J 6), 8.12 (2H, m), 9.10 (1H, s), 9.19 (1H, s).

Found:  $M^+$  320  $C_{19}H_{20}N_4O$  requires 320

## Example 7

### N-(3-Methoxyphenyl)-N'-(2-methyl-4-quinolinyl) urea

3-Methoxyphenyl isocyanate (0.83 ml, 6.3 mmol) in dry dichloromethane (30 ml) was added slowly to 4-aminoquinaldine (1g, 6.3 mmol) in dry toluene (30 ml) under a nitrogen atmosphere, and left to stir at room temperature for 19h. The precipitate which formed was filtered off, washed with cold 1:1 toluene/dichloromethane and dried *in vacuo*. The crude product was purified by recrystallization from ethanol to give the title compound (0.99g, 51%) as a white solid, m.p. 191-193°C.

NMR ( $D_6$ -DMSO)  $\delta$ : 2.6 (3H, s), 3.77 (3H, s), 6.62 (1H, m), 6.99 (1H, d, J 6), 7.22-7.28 (2H, m), 7.61 (1H, t, J 3), 7.72 (1H, t, J 3), 7.89 (1H, d, J 6), 8.14 (2H, m), 9.18 (1H, s), 9.35 (1H, s).

Found:  $M^+$  307  $C_{18}H_{17}N_3O_2$  requires 307

**Example 8****N-(3-Ethoxycarbonylphenyl)-N'-(2-methyl-4-quinolinyl) urea**

- 5 3-Ethoxycarbonylphenyl isocyanate (1g, 5.2 mmol) in dry dichloromethane (30 ml), was added slowly to 4-aminoquinaldine (0.83g, 5.2 mmol) in dry toluene (30 ml), under a nitrogen atmosphere, and left to stir at room temperature for 19h. The precipitate which formed was filtered off, washed with cold 1:1 toluene/dichloromethane and dried **in vacuo**. The crude product was chromatographed on silica gel, using dichloromethane as the eluant to give the title compound (0.78g, 43%) as white crystals, m.p. 165-170°C.

NMR (D<sub>6</sub>-DMSO) δ: 1.32 (3H, t, J 3), 2.6 (3H, s), 4.33 (2H, q, J 6), 7.48 (1H, t, J 6), 7.59-7.75 (4H, m), 7.9 (1H, d, J 6), 8.12 (2H, m), 8.22 (1H, s), 9.18 (1H, s), 9.57 (1H, s).

- 15 Found: M<sup>+</sup> 349 C<sub>20</sub>H<sub>19</sub>N<sub>3</sub>O<sub>3</sub> requires 349

**Example 9****N-(2-Methyl-4-quinolinyl)-N'-(3-trifluoromethylphenyl) urea**

- 20 α,α,α- Trifluoro-m-tolyl isocyanate (0.96 ml, 6.33 mmol) in dry dichloromethane (30 ml) was added slowly to 4-amino-quinaldine (1g, 6.33 mmol) in dry toluene (30 ml), under a nitrogen atmosphere. Following the procedure described in Example 4, gave the title compound (0.18g, 85%) as a white powder, m.p. 165-170°C.

- 25 NMR (D<sub>6</sub>-DMSO) δ: 2.58 (3H, s), 7.37 (1H, m), 7.55-7.61 (3H, m), 7.7 (1H, t J 6), 7.87 (1H, d, J 8), 8.10 (3H, m), 9.22 (1H, s), 9.60 (1H, s).

Found: M<sup>+</sup> 345 C<sub>18</sub>H<sub>14</sub>N<sub>3</sub>O F<sub>3</sub> requires 345

30 **Example 10**

**N-(Phenyl)-N'-(3-quinolinyl) urea**

Phenyl isocyanate (0.75 ml, 7 mmol) in dry dichloromethane (30 ml) was added slowly to 3 aminoquinoline (1g, 7 mmol) in dry toluene (30 ml) under a nitrogen atmosphere.

- 35 Following the procedure described in Example 7, gave the title compound (1.18g, 65%) as a white powder, m.p. 289-290°C.

NMR ( $D_6$ -DMSO)  $\delta$ : 7.0 (1H, t, J 6), 7.30 (2H, t, J 8), 7.49-7.61 (4H, m),  
7.88-7.97 (2H, m), 8.54 (1H, d, J 3), 8.82 (1H, d, J 3), 8.92  
(1H, s), 9.14 (1H, s).

- 5 Found: C, 72.78; H, 5.13; N, 15.98%  
 $C_{16}H_{13}N_3O$  requires C, 72.99; H, 4.98; N, 15.96%  
Found:  $M^+$  263  $C_{16}H_{13}N_3O$  requires 263

### Example 11

#### 10 N-(3-Chloro-4-methylphenyl)-N'-(3-pyridyl) urea hydrochloride

- Nicotinoyl azide (0.40g, 2.7 mmol) was stirred at reflux under nitrogen atmosphere in dry  
toluene (10 ml) for 1h, with gas evolution. The solution was cooled to ambient  
temperature, and 3-chloro-4-methylaniline (0.30 ml, 2.4 mmol) was added. The  
15 suspension so formed was stirred for 1h, when the solid was filtered off, washed with 1:1  
toluene/dichloromethane, and dried *in vacuo* at 70°C. This gave the free base of the title  
compound (0.64g, 85%) as a white solid.

- NMR ( $D_6$ -DMSO)  $\delta$ : 2.25 (3H, s), 7.23 (2H, m), 7.31 (1H, m), 7.70 (1H, s), 7.93  
20 (1H, m), 8.18 (1H, d, J 4), 8.59 (1H, d, J 2), 8.90 (2H, 2xs).

- N-(3-Chloro-4-methyl)-N'-(3-pyridyl) urea (0.55g, 2.1 mmol) was dissolved in hot ethanol  
(10 ml), and a solution of hydrogen chloride in ether (ca. 0.9M, 2.5 ml, ca. 2.3 mmol) was  
added. The suspension was cooled to ambient temperature, and the solid was filtered off,  
25 washed with cold ethanol, and dried *in vacuo* at 70°C. This gave the title compound  
(0.62g, 76%) as a white solid, m.p. 214.5-216°C.

- NMR ( $D_6$ -DMSO)  $\delta$ : 2.25 (3H, s), 7.25 (2H, m), 7.68 (1H, s), 7.92 (1H, dd, J 8,  
5), 8.33 (1H, d, J 8), 8.49 (1H, d), 9.07 (1H, s), 9.79 (1H, s),  
30 10.37 (1H, s).

Found: C, 51.4; H, 4.5; N, 14.5%  
 $C_{13}H_{12}ClN_3O \cdot HCl \cdot 0.25H_2O$  requires C, 51.6; H, 4.5; N, 13.9%  
Found:  $M^+$  261, 263.  $C_{13}H_{12}ClN_3O$  requires 261, 263.

### 35 Example 12

#### N-(3-Chloro-4-methylphenyl)-N'-(4-pyridyl) urea hydrochloride

3-Chloro-4-methylaniline (0.65 ml, 5.3 mmol) was stirred under nitrogen in

dichloromethane (15 ml) at 0°C as triethylamine (0.82 ml, 5.9 mmol) was added. To this mixture was then added phosgene in toluene solution (1.93M, 4.1 ml, 7.9 mmol). After stirring at 0°C for 0.5h, triethylamine (1.6 ml, 11.8 mmol) was added and, after a further 0.5h, 4-aminopyridine (0.50g, 5.3 mmol) was added. The mixture was stirred at ambient  
5 temperature for 16h, and then treated with sodium hydroxide solution (5M, ca. 1 ml). After 0.5h, it was diluted with water (50 ml) and dichloromethane (50 ml), and the precipitate was filtered off, washed with water, and dried *in vacuo* at 70°C. This gave the free base of the title compound (1.03g, 74%) as a white solid.

10 NMR (D<sub>6</sub>-DMSO) δ: 2.25 (3H, s), 7.23 (2H, m), 7.41 (2H, d, J 5), 7.67 (1H, s), 8.35 (2H, d, J 5) 8.99 (1H, s), 9.18 (1H, s).

N-(3-Chloro-4-methylphenyl)-N'-(4-pyridyl) urea (1.03g, 3.9 mmol) was treated with hydrogen chloride using the method of Example 11. This gave the title compound (0.95g,  
15 81%) as a white solid, m.p. 235-240°C (decomp.).

NMR (D<sub>6</sub>-DMSO) δ: 2.27 (3H, s), 7.28 (2H, m), 7.67 (1H, s), 7.89 (2H, d, J 6), 8.60 (2H, d, J 6), 10.09 (1H, s), 11.27 (1H, s).

Found: C, 50.6; H, 4.4; N, 13.7%

20 C<sub>13</sub>H<sub>12</sub>ClN<sub>3</sub>O.HCl . 0.59 H<sub>2</sub>O requires C, 50.6; H, 4.6; N, 13.6%

Found: M<sup>+</sup> 261, 263 C<sub>13</sub>H<sub>12</sub>ClN<sub>3</sub>O requires 261, 263.

### Example 13

#### N-(3-Pyridyl)-N'-(3-(trifluoromethyl)phenyl)urea

25

The title compound was prepared in 91% yield from 3-pyridyl isocyanate and 3-aminobenzotrifluoride; m.p. 180-184° C.

30 NMR (DMSO-d<sub>6</sub>) δ: 7.3 (2H, m), 7.55 (2H, m), 7.95 (1H, d, J 8), 8.0 (1H, s), 8.2 (1H, d, J 4), 8.6 (1H, d, J 2), 9.0 (1H, s), 9.2 (1H, s).

### Example 14

#### N-(3-Methylphenyl)-N'-(3-pyridyl)urea hydrochloride

35

The title compound was prepared in 87% yield from 3-aminopyridine and m-tolyl isocyanate, followed by salt formation with HCl; m.p. 182-183° C.

NMR (DMSO-d<sub>6</sub>) δ: 2.3 (3H, s), 6.85 (1H, d, J 7), 7.2 (1H, t, J 8), 7.3 (2H, m), 7.9 (1H, dd, J 8,5), 8.3 (1H, m), 8.5 (1H, d, J 5), 9.1 (1H, d, J 2), 9.5 (1H, s), 10.35 (1H, s).

**Example 15****N-(4-Chlorophenyl)-N'-(3-pyridyl)urea**

- 5 The title compound was prepared in 29% yield from 3-aminopyridine, 1,1'-carbonyldiimidazole and 4-chloroaniline; m.p. 207-209° C

NMR (DMSO-d<sub>6</sub>) δ: 7.3 (3H, m), 7.5 (2H, d, J 9), 7.95 (1H, m), 8.2 (1H, m), 8.6 (1H, d, J 2), 8.9 (1H, s), 9.0 (1H, s)

10

**Example 16****N-(3-Chlorophenyl)-N'-(3-pyridyl)urea**

- 15 The title compound was prepared in 86% yield from 3-aminopyridine and 3-chlorophenyl isocyanate; m.p. 185-187° C

NMR (DMSO-d<sub>6</sub>) δ: 7.0 (1H, m), 7.3 (3H, m), 7.7 (1H, s), 7.95 (1H, m), 8.2 (1H, m), 8.6 (1H, d, J 2), 8.95 (1H, s), 9.05 (1H, s)

20 **Example 17**

**N-(3-Hydroxyphenyl)-N'-(2-methyl-4-quinolinyl)urea**

- 25 The title compound was prepared in 19% yield from 4-amino-2-methylquinoline, 1,1'-carbonyldiimidazole and 3-aminophenol; m.p. 224-225° C

NMR (DMSO-d<sub>6</sub>) δ: 2.6 (3H, s), 6.45 (1H, m), 6.9 (1H, d, J 7), 7.1 (2H, m), 7.6 (1H, t, J 7), 7.7 (1H, t, J 7), 7.9 (1H, d, J 7), 8.15 (2H, m), 9.2 (1H, b), 9.3 (1H, s), 9.45 (1H, s)

30 **Example 18**

**N-(3-Bromophenyl)-N'-(3-pyridyl)urea**

- The title compound was prepared in 75% yield from 3-bromopyridine and 3-pyridyl isocyanate; m.p. 190-193° C.

35 NMR (DMSO-d<sub>6</sub>) δ: 7.10-7.40 (4H, m), 7.86 (1H, s), 7.94 (1H, m), 8.22 (1H, d, J=5Hz), 8.62 (1H, d, J=2Hz), 8.93 (1H, s), 9.02 (1H, s).

**Example 19****N-(3,4-Dichlorophenyl)-N'-(3-pyridyl)urea**

40

The title compound was prepared in 65% yield from 3,4-dichloroaniline and 3-pyridyl isocyanate; m.p. 206° C-210° C.

45 NMR (DMSO-D<sub>6</sub>) δ: 7.25-7.42 (2H, m), 7.50 (1H, d, J=7Hz), 7.83-7.90 (2H, m), 8.23 (1H, d, J=3Hz), 8.62 (1H, d, J=1Hz), 8.98 (1H, s), 9.23 (1H, s)



**Example 20****N-(3-Fluoro-4-methylphenyl)-N'-(3-pyridyl)urea**

5 The title compound was prepared in 85% yield from 3-fluoro-4-methylaniline and 3-pyridyl isocyanate; m.p. 190-191° C.

NMR (DMSO-D<sub>6</sub>) δ: 7.02-7.48 (4H, m), 7.94 (1H, m), 8.19 (1H, m), 8.59 (1H, m), 8.87 (1H, s), 8.92 (1H, s)

10

**Example 21****N-(4-Ethoxycarbonylphenyl)-N'-(3-pyridyl)urea**

15 The title compound was prepared in 83% yield from ethyl 4-aminobenzoate and 3-pyridyl isocyanate; m.p. 156-160° C

NMR (DMSO-D<sub>6</sub>) δ: 1.32 (3H, t, J=7.5Hz), 4.30 (2H, q, J=7.5Hz), 7.34 (1H, dd, J=7Hz & 4Hz), 7.60 (2H, m), 7.86-8.02 (3H, m), 8.21 (1H, m), 8.63 (1H, m), 8.96 (1H, s), 9.24 (1H, s)

20

**Example 22****N-(3-Chloro-4-methoxycarbonylphenyl)-N'-(3-pyridyl)urea**

25 The title compound was prepared in 30% yield from methyl 4-amino-2-chlorobenzoate and 3-pyridyl isocyanate m.p. 170-171° C

NMR (DMSO-D<sub>6</sub>) δ: 3.82 (3H, s), 7.30 (2H, m), 7.78-8.00 (3H, m), 8.25 (1H, m), 8.64 (1H, m), 9.08 (1H, s), 9.39 (1H, s)

30

**Example 23****N-(3-Bromo-4-methylphenyl)-N'-(3-pyridyl)urea**

The title compound was prepared in 61% yield from 3-bromo-4-methylaniline and 3-pyridyl isocyanate; m.p. 168-171° C

35

NMR (DMSO-D<sub>6</sub>) δ: 2.28 (3H, s), 7.21-7.39 (3H, m), 7.83-8.00 (2H, m), 8.20 (1H, m), 8.61 (1H, m), 8.89 (2H, m)

**Example 24**

40 **N-(3-Chloro-4-cyanophenyl)-N'-(3-pyridyl)urea**

The title compound was prepared in 22% yield from 4-amino-2-chlorobenzonitrile and 3-pyridyl isocyanate; m.p. 262-264° C

NMR (DMSO-D<sub>6</sub>)  $\delta$ : 7.28-7.56 (2H, m), 7.80-8.06 (3H, m), 8.26 (1H, m), 8.64 (1H, s), 9.17 (1H, s), 9.54 (1H, s)

**Example 25**

5 **N-(4-Nitro-3-trifluoromethylphenyl)-N'-(3-pyridyl)urea**

The title compound was prepared in 31% yield from 4-nitro-3-trifluoromethylaniline and 3-pyridyl isocyanate; m.p. 214-216° C

10 NMR (DMSO-D<sub>6</sub>)  $\delta$ : 7.37 (1H, dd, J=7Hz & 4Hz), 7.87 (1H, m, J=7Hz), 7.97 (1H, m, J=7Hz), 8.14-8.29 (3H, m), 8.67 (1H, m), 9.22 (1H, s), 9.81 (1H, s)

**Example 26**

15 **N-(4-Chloro-3-trifluoromethylphenyl)-N'-(3-pyridyl)urea**

The title compound was prepared in 48% yield from 4-chloro-3-trifluoromethylaniline and 3-pyridyl isocyanate; m.p. 196-199° C.

20 NMR (DMSO-D<sub>6</sub>)  $\delta$ : 7.33 (1H, dd, J=7Hz & 4Hz), 7.59-7.71 (2H, m), 7.95 (1H, m), 8.10 (1H, m), 8.22 (1H, m), 8.63 (1H, m), 9.04 (1H, s), 9.32 (1H, s)

**Example 27**

**N-(3-Chloro-4-carboxyphenyl)-N'-(3-pyridyl)urea**

25 The title compound was prepared in 86% yield from 4-amino-2-chlorobenzoic acid and 3-pyridyl isocyanate; m.p. 170-175° C

30 NMR (DMSO-D<sub>6</sub>)  $\delta$ : 7.41 (2H, m), 7.76-7.88 (2H, m), 7.99 (1H, d, J=7Hz), 8.25 (1H, br s), 8.68 (1H, br s), 9.13 (1H, s), 9.37 (1H, s)

**Example 28**

**N-(2-Methoxy-4-trifluoromethylphenyl)-N'-(3-pyridyl)urea**

35 The title compound was prepared in 91% yield from 2-methoxy-4-trifluoromethyl-aniline and 3-pyridyl isocyanate; m.p. 210° C

NMR (DMSO-D<sub>6</sub>)  $\delta$ : 4.00 (3H, s), 7.16-7.45 (3H, m), 7.98 (1H, m, J=7Hz), 8.23 (1H, m), 8.48-8.74 (3H, m), 9.60 (1H, s)

40 **Example 29**

**N-(2,3-Dichlorophenyl)-N'-(2-methyl-4-quinoliny)urea**

The title compound was prepared in 22% yield from 2,3-dichloroaniline and 2-methyl-4-quinoliny isocyanate; m.p. 125-127° C

45

NMR (DMSO-D<sub>6</sub>)  $\delta$ : 2.62 (3H, s), 7.34-7.46 (2H, m), 7.63 (1H, t, J=7Hz), 7.76 (1H, t, J=7Hz), 7.94 (1H, t, J=7Hz), 8.12-8.31 (3H, m), 9.27 (1H, s), 9.83 (1H, s)

### Example 30

#### 5 N-(3-Chloro-4-ethylphenyl)-N'-(3-pyridyl)urea

The title compound was prepared in 85% yield from 3-chloro-4-ethylaniline & 3-pyridyl isocyanate; m.p. 193-196° C.

10 NMR (DMSO-d<sub>6</sub>)  $\delta$ : 1.16 (3H, t, J=5Hz), 2.64 (2H, q, J=5Hz), 7.20-7.40 (3H, m), 7.67 (1H, s), 7.94 (1H, m), 8.20 (1H, d, J=2Hz), 8.60 (1H, d, J=0-1Hz), 8.90 (2H, d, J=5Hz).

### Example 31

#### 15 N-(3-Chloro-4-propylphenyl)-N'-(3-pyridyl)urea

The title compound was prepared in 78% yield from 3-Chloro-4-propylaniline & 3-pyridyl isocyanate; m.p. 184-186° C

20 NMR (DMSO-D<sub>6</sub>)  $\delta$ : 0.91 (3H, t, J=5Hz), 1.56 (2H, q, J=5Hz), 2.60 (2H, t, J=5Hz), 7.20-7.35 (3H, m), 7.68 (1H, s), 7.94 (1H, m), 8.19 (1H, d, J=2Hz), 8.59 (1H, d, J=0-1Hz), 8.92 (2H, d, J=5Hz).

### Example 32

#### 25 N-(3-Chloro-4-tert-butylphenyl)-N'-(3-pyridyl)urea

The title compound was prepared in 73% yield from 3-chloro-4-tert-butylniline & 3-pyridyl isocyanate; m.p. 190° C-193° C.

30 NMR (DMSO-D<sub>6</sub>)  $\delta$ : 1.42 (9H, s), 7.20-7.40 (3H, m), 7.66 (1H, d, J=2Hz), 7.93 (1H, m), 8.19 (1H, d, J=5Hz), 8.60 (1H, d, J=2Hz), 8.90 (2H, d, J=11Hz)

### Example 33

#### N-(3-Hydroxy-4-(methoxycarbonyl)phenyl)-N'-(3-pyridyl)urea

35 N-(3-Hydroxy-4-carboxyphenyl)-N'-(3-pyridyl)urea was prepared in 69% yield from 4-aminosalicylic acid and 3-pyridyl isocyanate in DMF/toluene. This material (0.37g, 1.4 mmol) was then stirred in methanol (20 ml) as thionyl chloride (2 ml) was cautiously added. The suspension was stirred at reflux under argon for 2 days, and evaporated to dryness. The residue was suspended in saturated sodium hydrogen carbonate solution, and  
40 the solid was filtered off, washed with water, dried, and recrystallised from ethanol/petroleum ether (b.p. 60-80° C), giving the title compound (0.16g, 41%) as a white solid. m.p. 199-200° C.

NMR (DMSO  $d_6$ )  $\delta$  :

3.88 (3H, s), 6.98 (1H, dd, J 8, 2), 7.27 (1H, d, J 2), 7.34 (1H, dd, J 8, 5), 7.73 (1H, d, J 9), 7.96 (1H, m), 8.24 (1H, d, J 4), 8.63 (1H, d, J 2), 9.04 (1H, s), 9.27 (1H, s), 10.69 (1H, s).

5

**Pharmacological data****[<sup>3</sup>H]-mesulergine binding to rat 5-HT<sub>2C</sub> clones expressed in 293 cells in vitro**

Evidence from the literature suggests that 5-HT<sub>2C</sub> antagonists may have a number of therapeutic indications including the treatment of anxiety, migraine, depression, feeding disorders and obsessive compulsion disorders. (Curzon and Kennett, 1990; Fozard and Gray, 1989) and Alzheimer's Disease (Lawlor, 1989, J. Arch. Gen. Psychiat. Vol. 46 p.542).

10 The affinity of test drugs for the 5-HT<sub>2C</sub> binding site can be determined by assessing their ability to displace [<sup>3</sup>H]-mesulergine from 5-HT<sub>2C</sub> clones expressed in 293 cells (Julius *et al.*, 1988). The method employed was similar to that of Pazos *et al.*, 1984.

15 The cells suspension (50ml) was incubated with [<sup>3</sup>H]-mesulergine (0.5nM) in Tris HCl buffer (pH 7.4) at 37°C for 30 minutes. Non-specific binding was measured in the presence of mianserin (10<sup>-6</sup>M). Ten concentrations of test drug (3 x 10<sup>-9</sup> to 10<sup>-4</sup>M final concentration) were added in a volume of 50ml. The total assay volume was 500ml. Incubation was stopped by rapid filtration using a Brandel cell harvester and radioactivity measured by scintillation counting. The IC<sub>50</sub> values were determined using  
20 a four parameter logistic program (DeLean 1978) and the pK<sub>i</sub> (the negative logarithm of the inhibition constant) calculated from the Cheng Prusoff equation where:

$$K_i = \frac{IC_{50}}{1 + \frac{C}{K_d}}$$

25

K<sub>i</sub> = inhibition constant.  
30 C = concentration of [<sup>3</sup>H]-mesulergine  
K<sub>d</sub> = Affinity of mesulergine for 5-HT<sub>1C</sub> binding sites.

Curzon, G.A. and Kennett, G.A. (1990). TIPS, Vol. 11, 181-182.  
Fozard, J.R. and Gray, J.A. (1989). TIPS, Vol. 10, 307-309.  
35 Pazos, A. *et al.* (1984). Eur. J. Pharmacol., 106, 531-538.  
Julius *et al.* (1988) Science 241, 558-564  
DeLean A, Munson P.J., Rodbaud D (1978) Am. J. Physiol 235, E97-E102.

## Results

The compound of Example 7 has a pKi of 8.28.

The compound of Example 11 has a pKi of 7.79.

### 5 Reversal of MCPP-induced Hypolocomotion

Administration of m-(chlorophenyl)piperazine (mCPP) to rats induces hypolocomotion (Kennett and Curzon 1988, Luckie *et al.* 1989) as seen with the related drug 1-(m-trifluoromethylphenyl)piperazine (TFMPP) (Lucki and Frazer 1982, Kennett and Curzon 1988). This effect was blocked by the non specific

- 10 5-HT<sub>2C</sub>/5-HT<sub>2A</sub> receptor antagonists mianserin, cyproheptadine and metergoline and perhaps by mesulergine. It was not blocked by the 5-HT<sub>2</sub> receptor antagonists ketanserin and ritanserin at relevant doses (Kennett and Curzon 1991) nor by antagonists of 5-HT<sub>1A</sub>, 5-HT<sub>1B</sub>, 5-HT<sub>3</sub>,  $\alpha_2$  adrenoceptors or dopamine D<sub>2</sub> receptors. The effect of mCPP is therefore considered to be mediated by 5-HT<sub>2C</sub> receptors (Kennett and Curzon 1988) as
- 15 confirmed by subsequent studies (Lucki *et al.*, 1989). Since mCPP causes hypolocomotion when infused into the cerebral ventricles this effect is probably centrally mediated (Kennett and Curzon 1988).

- mCPP-induced hypolocomotion was measured in automated locomotion cages of
- 20 dimensions 56 cm long x 16½ cm wide x 25 cm high and made of black perspex. Two photobeams traversed the width of the cages at either end at ground level. Sequential breaking of these beams allowed the measurement of cage transits.

- Male Sprague Dawley rats (200-250g) (Charles River) were housed in groups of six. They
- 25 were given drugs orally 1h pretest and 40 mins later mCPP (7 mg/kg i.p.). After a further 20 min they were placed in individual automated cages in groups of four under red light in an adjacent room. After 10 min the test was terminated. Reversal of mCPP-induced hypolocomotion was considered as evidence of *in vivo* central 5-HT<sub>2C</sub> receptor antagonist properties.

30

Kennett, G.A., Curzon, G., (1988). Brit. J. Pharmacol. **94**, 137-147.

Kennet G.A., Curzon, G., (1991). Brit.J. Pharmacol. **103**, 2016-2020.

Lucki, I., Frazer, A., (1982) Am. Soc. Neurosci. 8(abstr.), 101.

Lucki, I., Ward, M.R., Frazer, A., (1989). J.Pharmacol. Exp. Therap. **249**, 155-164.


35

## Result

The compound of Example 11 had an ID<sub>50</sub> of 78 mg/kg p.o.

## CLAIMS

1. Use of a compound of formula (I) or a salt thereof:



5

wherein:

- P represents a quinoline or isoquinoline residue or a 5- or 6-membered aromatic heterocyclic ring containing up to three heteroatoms selected from nitrogen, oxygen or sulphur:

- $R^1$  is hydrogen,  $C_{1-6}$  alkyl, halogen,  $NR^5R^6$  or  $OR^7$  where  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen or  $C_{1-6}$  alkyl;  
 $R^2$  and  $R^3$  are independently hydrogen or  $C_{1-6}$  alkyl;  
 $R^4$  is hydrogen,  $C_{1-6}$  alkyl,  $CF_3$ , nitro, cyano, acyl, halogen,  $NR^5R^6$ ,  $OR^7$  or  $CO_2R^7$  where  $R^5$ ,  $R^6$  and  $R^7$  are independently hydrogen or  $C_{1-6}$  alkyl as defined for  $R^1$ ; and  $n$  is 1, 2 or 3,  
 in the manufacture of a medicament for the treatment or prophylaxis of CNS disorders.

2. Use according to claim 1 in which P is pyridyl or quinolyl.
3. Use according to claim 1 or 2 in which  $R^1$  is hydrogen or methyl.
4. Use according to any one of claims 1 to 3 in which  $R^2$  and  $R^3$  are hydrogen.
5. Use according to any one of claims 1 to 4 in which P is pyridyl or quinolyl.
6. Use according to claim 1 in which the compound of formula (I) is selected from:  
 N-(Phenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(6-Quinoliny)l)-N'-(3-trifluoromethylphenyl) urea,  
 N-(3-Dimethylaminophenyl)-N'-(6-quinoliny)l urea,  
 N-(Phenyl)-N'-(6-quinoliny)l urea,  
 N-(4-Methoxyphenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(3-Dimethylaminophenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(3-Methoxyphenyl)-N'-(2-methyl-4-quinoliny)l urea,


- N-(3-Ethoxycarbonylphenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(2-Methyl-4-quinoliny)l-N'-(3-trifluoromethylphenyl) urea ,  
 N-(Phenyl)-N'-(3-quinoliny)l urea,  
 N-(3-Chloro-4-methylphenyl)-N'-(3-pyridyl) urea,  
 5 N-(3-Chloro-4-methylphenyl)-N'-(4-pyridyl) urea,  
 N-(3-Pyridyl)-N'-(3-(trifluoromethyl)phenyl)urea,  
 N-(3-Methylphenyl)-N'-(3-pyridyl)urea,  
 N-(4-Chlorophenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chlorophenyl)-N'-(3-pyridyl)urea,  
 10 N-(3-Hydroxyphenyl)-N'-(2-methyl-4-quinoliny)lurea,  
 N-(3-Bromophenyl)-N'-(3-pyridyl)urea,  
 N-(3,4-Dichlorophenyl)-N'-(3-pyridyl)urea,  
 N-(3-Fluoro-4-methylphenyl)-N'-(3-pyridyl)urea,  
 N-(4-Ethoxycarbonylphenyl)-N'-(3-pyridyl)urea,  
 15 N-(3-Chloro-4-methoxycarbonylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Bromo-4-methylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-cyanophenyl)-N'-(3-pyridyl)urea,  
 N-(4-Nitro-3-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
 N-(4-Chloro-3-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
 20 N-(3-Chloro-4-carboxyphenyl)-N'-(3-pyridyl)urea,  
 N-(2-Methoxy-4-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-ethylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-propylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-tert-butylphenyl)-N'-(3-pyridyl)urea,  
 25 N-(3-Hydroxy-4-(methoxycarbonyl)phenyl)-N'-(3-pyridyl)urea  
 or a pharmaceutically acceptable salt thereof.

7. A compound of formula (I) which is:  
 N-(Phenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 30 N-(6-Quinoliny)l-N'-(3-trifluoromethylphenyl) urea,  
 N-(3-Dimethylaminophenyl)-N'-(6-quinoliny)l urea,  
 N-(Phenyl)-N'-(6-quinoliny)l urea,  
 N-(4-Methoxyphenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(3-Dimethylaminophenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 35 N-(3-Methoxyphenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(3-Ethoxycarbonylphenyl)-N'-(2-methyl-4-quinoliny)l urea,  
 N-(2-Methyl-4-quinoliny)l-N'-(3-trifluoromethylphenyl) urea ,  
 N-(Phenyl)-N'-(3-quinoliny)l urea,




- N-(3-Chloro-4-methylphenyl)-N'-(3-pyridyl) urea,  
 N-(3-Chloro-4-methylphenyl)-N'-(4-pyridyl) urea,  
 N-(3-Pyridyl)-N'-(3-(trifluoromethyl)phenyl)urea,  
 N-(3-Methylphenyl)-N'-(3-pyridyl)urea,  
 5 N-(4-Chlorophenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chlorophenyl)-N'-(3-pyridyl)urea,  
 N-(3-Hydroxyphenyl)-N'-(2-methyl-4-quinoliny)urea,  
 N-(3-Bromophenyl)-N'-(3-pyridyl)urea,  
 N-(3,4-Dichlorophenyl)-N'-(3-pyridyl)urea,  
 10 N-(3-Fluoro-4-methylphenyl)-N'-(3-pyridyl)urea,  
 N-(4-Ethoxycarbonylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-methoxycarbonylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Bromo-4-methylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-cyanophenyl)-N'-(3-pyridyl)urea,  
 15 N-(4-Nitro-3-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
 N-(4-Chloro-3-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-carboxyphenyl)-N'-(3-pyridyl)urea,  
 N-(2-Methoxy-4-trifluoromethylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-ethylphenyl)-N'-(3-pyridyl)urea,  
 20 N-(3-Chloro-4-propylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Chloro-4-tert-butylphenyl)-N'-(3-pyridyl)urea,  
 N-(3-Hydroxy-4-(methoxycarbonyl)phenyl)-N'-(3-pyridyl)urea  
 or a pharmaceutically acceptable salt thereof.
- 25 8. A process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt thereof which comprises:

the coupling of a compound of formula (II);



with a compound of formula (III);



(III)

wherein P is as defined in relation to formula (I), A and B contain the appropriate

functional group(s) necessary to form the moiety,  $-NR^{2'}CONR^{3'}$  when coupled, the variables  $R^{1'}$ ,  $R^{2'}$ ,  $R^{3'}$ , and  $R^{4'}$  are  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  respectively, as defined in formula (I), or groups convertible thereto, and thereafter optionally and as necessary and in any appropriate order, converting any  $R^{1'}$ ,  $R^{2'}$ ,  $R^{3'}$  and  $R^{4'}$ , when other than  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$  respectively to  $R^1$ ,  $R^2$ ,  $R^3$  and  $R^4$ , interconverting  $R^1$ ,  $R^2$ ,  $R^3$ , and  $R^4$  and forming a pharmaceutically acceptable salt thereof.

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 94/00189

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C07D213/75 C07D215/38 C07D215/46 A61K31/47 A61K31/44

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,92 05170 (BEECHAM GROUP PLC) 2 April 1992 cited in the application see the whole document ---	1-8
P,Y	WO,A,93 18028 (SMITHKLINE BEECHAM PLC) 16 September 1993 *see whole document, especially definition of P* ---	1-8
A	JOURNAL OF MEDICINAL CHEMISTRY vol. 29, no. 11, 1986, WASHINGTON US pages 2415 - 2418 P. FLUDZINSKI ET AL '2,3-dialkyl(dimethylamino)indoles: Interaction with 5HT <sub>1</sub> , 5HT <sub>2</sub> and rat stomach fundal serotonin receptors.' *see especially compound number 10* --- -/--	1-8

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 April 1994

Date of mailing of the international search report

09-05-94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Scruton-Evans, I

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 94/00189

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 515 684 (CHUGAI SEIYAKU KABUSHIKI KAISHA) 2 December 1992 *see compounds of formula I wherein R is (I)-2, and their process of preparation on page 8*	8
A	---	1-7
A	PATENT ABSTRACTS OF JAPAN vol. 16, no. 473 (C-0991) 2 October 1992 & JP,A,04 173 701 (HOKKO CHEM IND CO LTD) 22 June 1992 see abstract	1-7
X	---	
X	US,A,4 880 817 (R.M.KANOJIA ET AL) 14 November 1989 *see column 6, lines 64-68 and column 7, lines 1-14*	8
A	---	1-7
P,A	---	
P,A	EP,A,0 540 854 (SANSHO SEIYAKU CO. LTD.) 12 May 1993 *see compounds of formula 2 and definitions of R1 and R2 on page 6*	1-7
A	---	
A	EP,A,0 354 994 (TAKEDA CHEMICAL INDUSTRIES LTD) 21 February 1990 -----	1-8

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 94/00189

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9205170	02-04-92	AU-B- 642041 AU-A- 8503891 CA-A- 2091246 EP-A- 0550507 JP-T- 6500551	07-10-93 15-04-92 14-03-92 14-07-93 20-01-94
WO-A-9318028	16-09-93	NONE	
EP-A-0515684	02-12-92	AU-A- 7317391 WO-A- 9111994	03-09-91 22-08-91
US-A-4880817	14-11-89	NONE	
EP-A-0540854	12-05-93	AU-A- 2281792 CA-A- 2077850	11-03-93 11-03-93
EP-A-0354994	21-02-90	AU-B- 616542 AU-A- 3802589 US-A- 5254565 JP-A- 3007259	31-10-91 18-01-90 19-10-93 14-01-91